


Spike Protein Distribution



Discovered after COVID-19 infection



Discovered after COVID-19 vaccination



Note: The times listed for spike protein presence in different organs are based on specific and separate studies, and may not be comprehensive or comparable between times listed.

Persistence of SARS-CoV-2 S1 Protein in CD16+ Monocytes in Post-Acute Sequelae of COVID-19 (PASC) up to 15 Months Post-Infection, *Frontiers in Immunology*
 Cell death and pathological findings of the spleen in COVID-19 patients, *Elsevier Public Health Emergency Collection*
 The SARS-CoV-2 Spike protein disrupts human cardiac pericytes function through CD147 receptor-mediated signaling: a potential non-infective mechanism of COVID-19 microvascular disease, *Clinical Science*
 The central role of the nasal microenvironment in the transmission, modulation, and clinical progression of SARS-CoV-2 infection, *Mucosal Immunology*
 Mucosal Immunology, *Nat Med*
 Oral SARS-CoV-2 Inoculation Causes Nasal Viral Infection Leading to Olfactory Bulb Infection: An Experimental Study, *Frontiers in Cellular and Infection Microbiology*
 Evidence of SARS-CoV-2 in nasal brushings and olfactory mucosa biopsies of COVID-19 patients, *PLOS ONE*
 SARS-CoV-2 infection of the oral cavity and saliva, *Nature Medicine*
 Circulating SARS-CoV-2 Vaccine Antigen Detected in the Plasma of mRNA-1273 Vaccine Recipients, *Oxford University Press*
 SARS-CoV-2 infects adipose tissue in a fat depot- and viral lineage-dependent manner, *Nature Communications*
 The spike protein of SARS-CoV-2 induces endothelial inflammation through integrin $\alpha 5 \beta 1$ and NF- κB signaling, *Journal of Biological Chemistry*
 The S1 protein of SARS-CoV-2 crosses the blood-brain barrier in mice, *Nature Neuroscience*
 SARS-CoV-2 S1 Protein Induces Endolysosome Dysfunction and Neuritic Dystrophy, *Frontiers in Cellular Neuroscience*
 SARS-CoV-2 spike protein causes cardiovascular disease independent of viral infection, *Clinical Science*
 Detection of SARS-CoV-2 antigens in thyroid gland showing histopathological features of subacute thyroiditis, *European Thyroid Journal*
 Covid vaccine induced subacute thyroiditis, *Organizace Mafialda Saude*
 SARS-CoV-2 in the pancreas and the impaired islet function in COVID-19 patients, *Emerging Microbes & Infections: CDC Morbidity and Mortality Weekly Report*
 Nephrotic Syndrome After Vaccination Against COVID-19: Three New Cases From Germany, *Deutscher Arztekongress*
 Evidence for SARS-CoV-2 Spike Protein in the Urine of COVID-19 Patients, *Kidney360*
 COVID-19 targets human adrenal glands, *Elsevier Public Health Emergency Collection*

Adrenal tropism of SARS-CoV-2 and adrenal findings in a post-mortem case series of patients with severe fatal COVID-19, *Nature Communications*
 Pfizer SARS-CoV-2 mRNA Vaccine (BNT162z, PF-07302048) pharmacokinetic study, *Presumed SARS-CoV-2 Viral Particles in the Human Retina of Patients With COVID-19*, *JAMA Ophthalmology*
 Retinal findings in patients with COVID-19: Results from the SERPICO-19 study, *Lancet Discovery Science*
 COVID-19 chibblain-like lesion: immunohistochemical demonstration of SARS-CoV-2 spike protein in blood vessel endothelium and sweat gland epithelium in a polymerase chain reaction-negative patient, *Oxford University Press*
 SARS-CoV-2 spike protein is present in both endothelial and eccrine cells of a chibblain-like skin lesion, *Journal of the European Academy of Dermatology & Venereology*
 Post-COVID-19 menstrual abnormalities and infertility: Repercussions of the pandemic, *Journal of Education and Health Promotion*
 COVID-19 and the surge in Decidual Cast Shedding, *The Gazette of Medical Sciences*
 SARS-CoV-2 Infection of Human Ovarian Cells: A Potential Negative Impact on Female Fertility, *Cells*
 Coronavirus: Why Men are More Vulnerable to Covid-19 Than Women?, *Nature Public Health Emergency Collection*
 SARS-CoV-2 infects, replicates, elevates angiotensin II and activates immune cells in human testes, pre-print
 Pathological and molecular examinations of postmortem testis biopsies reveal SARS-CoV-2 infection in the testis and spermatogenesis damages in COVID-19 patients, *Cellular Molecular Immunology*
 Semen impairment and occurrence of SARS-CoV-2 virus in semen after recovery from COVID-19, *Oxford University Press*
 CDC Dispatch SARS-CoV-2 RNA Shedding in Semen and Oligospermia of Patient with Severe Coronavirus Disease 11 Weeks after Infection
 Covid-19 vaccination BNT162b2 temporarily impairs semen concentration and total motile count among semen donors, *Andrology*
 Effects of COVID-19 vaccine on semen parameters, *Nature Reviews*